

What is an Antitoxin?

- Antitoxins are NON-toxic and globally unrestricted **first aid**.
- Antitoxins are biological, without animal derivatives.
- Antitoxins are concentrates that can be used as food supplementation.
- Antitoxins help in the management of potentially dangerous foods.
- Antitoxins can prevent poison processes and proven by blood test document.

\$nake Antidote is a biological plant-based detox supplement made of bitter iron concentrate which doesn't contain any bio-proteins or active salts or natron. It is not administered with a dangerous injection needle. It is used with speed, orally.

How does classical antitoxin work:

\$nake Antidote is a dark biological liquid tannin iron which functions by staining the blood so as to reduce the overt **toxic level** of viro-proteins in the blood that produce **hyper insulin volume** (HIV-complex) in sicknesses such as:

- **Malaria**
- **Leukaemia**
- **Hepatitis**
- **Sickle-cell Anaemia**
- **Iron deficiency**

Hyper insulin volume (HIV-complex) is caused by unbalanced white salty body fats that destroy body cells, and is the opposite of **hypo insulin volume** (HIV-simplex). - "*Hyper*" means high, and "*Hypo*" means low.

\$nake Antidote converts toxins into a non-toxic waste material, which will leave the body by following the laws of gravity in iron's magnetism to the earth physics. - Expulsion. This purger cannot be used to treat **hypo insulin volume** (HIV-simplex / insulin deficiency) which is the opposite of **hyper insulin volume** (HIV-complex / insulin overdose). **Hypo insulin volume** (HIV-simplex / insulin deficiency) is an insulin deficit leading to acidic insulin deficiency through **hyper peptide volume** (HPV) – a sulfurisation that leads to **acid insulin deficiency sulfurisation** (AIDS-complex / acid insulin deficiency sulfurisation) in chemical police forensics.

Classic biological antitoxins are safe to use when...

Bit by a poisonous animal, or when diets, lifestyle, persons and places have become toxic through bacterial imbalance, decay, poison, gases, smells or drug traffic. Antitoxins are taken in recommended small to large doses to treat diverse toxic symptoms in humans or animals.

The following pages offer quick speedy explanation on when, why and how to use classic antitoxin. Study this long in advance of any emergency.

**FIRST AID
USE AT A GLANCE**

ORAL SNAKE ANTIDOTE

Amount per person average large poisoning = **1 unit** = 2 bottles, with speed!

APPLICATION:

1. Contravention of venomous bites from toxic snakes, spiders, scorpions, stingray, jellyfish, colour-frogs, venomous marine life.
2. The treatment of the uncertainty about if a bite was toxic or not.
3. Treatment against toxic gases of war and food poisoning.
4. Treatment of palliative drug addiction (Cocaine, Heroine, Tranquillizers etc.)
5. Treatment of blood-documented **Hyper-Insulin-Volume** (HIV-complex) levels such as in:
 - Malaria
 - Leukaemia
 - Hepatitis
 - Sickle-cell Anaemia
 - Iron deficiency
6. As a diuretic treating filaria and thrombosis.
7. For the treatment of obesity, overweight and cholesterol.
8. To treat overweight diabetics along a tailored diet plan.
9. In mild doses coupled with a protein source to raise low sexual libido.
10. In small doses to promote memory, keen concentration, and relaxation.
11. To bitter the blood occasionally.
12. As a purger to treat constipation.

YOUR BASIC LEGAL RIGHT for FIRST AID SELF DEFENCE

Humans and pets take dangerous risks all the time and some doctors give **first-aid classes** to help you to help yourself and others. Waiting for a paramedic to arrive on site can waste precious lifetime **when you live far away** and you or a loved one has been seriously poisoned. Saving livestock saves from the loss of livelihood and income. This simple biological product is intended as a high-speed self-aid detox for vulnerable helpless people who may have difficult access to medical assistance in a lonely place of danger. It is better to have biological antitoxin safety, than no help arriving too late. **It's YOUR life!**

The ancient Egyptian and Greek snake antidote **symbol** of a snake on a mono-pole is displayed on the dollar and every pharmacy sign in the world, yet these antitoxins aren't even in them. Why? The exorbitant cost of less effective US- and EU-made snake antidote ranges between 370 Euros

to 11.000 \$ for just one single vial of 10ml representing 1 part of a 5-vial treatment at 55.000 \$ – a price that rural poor working classes of the world and BRICS cannot afford. These Western non-biological antidotes are derived from animal plasma and aren't PETA conform. In the ancient world this \$ymbol of money did not come at such a deadly price. Moreover, animal blood was **not** used to make them.

Now supposing, that an inviting meal or drink has been contaminated, poisoned or even drugged; isn't it highly unwise to have absolutely nothing to aid your personal self-defence and immunity? A cheap concentrated oral antidote is better than regretting the lack. **Most people have justified doubts about recognizing the difference between a non-venomous snake and a venomous one, when surprised by a bite.** Often the elusive snake retracts so quickly into hiding and even escapes the scene before we can even identify it's specimen. It is best not to waste time guessing which snakelike person or animal has hit, but to be prepared in advance. If your child or dog is dizzy or limping after playing in the tropical garden it is best to quickly search for a scorpion sting or reptile mark and quickly administer the emergency solution. **It's your basic right!** Antitoxins are **non-toxic HELP!** - Life is short, as it stands.

Safety Necessity:

This biological antitoxin is **broad spectrum**. There are **three toxic poison groups** that loosely apply to most existing poisons on the planet whether acute, chronic or fatal, which will decide which detox process is correct for your defence:

1. **Haemotoxic** = causing simple swelling or chronic carcinogenesis or deadly necrosis.
2. **Neurotoxic** = causing disorientation or sudden comatosis or death from overdose.
3. **Cytotoxic** = causing minor infection to instant gangrene decay and abrupt death.

Ascertain what type of poison risk lurks in your destination or backyard first before setting out. Depending on the severity, the first two groups - **haemotoxic** and **neurotoxic** – can become deadly, but easier to treat. They are **palliative** poisons that require simple but speedy classic antitoxin application called *“open bottle – drink it – keep still!”*. However, **Cytotoxic**, means combating highly sulphurous agents that cause infections and if left untreated can advance fatally. High cytotoxicity can lead to sudden asphyxiation, cell death and sulphurous decay that may be harder to treat, because inhaling a chlorine corona gas - **can choke you in an instant**. Getting hit by a cytotoxic serpent anywhere other than the legs, can speed extremely chronic sulphurs to your heart, head, causing a brain stroke with excruciating tissue and neural damage. Therefore, **Cytotoxins** require that you **take the extra trouble** of initially swallowing steroids (i.e. Calcium and/or white BIOTIN pills), rather than an iron antitoxin per se. When taking a unit of iron antidote **before** you set out into a zone known for cytotoxic snakes or sulphurous chlorine gases of war, consider also having calcium or biotin pills along. The steroids will spread with a meal in the body to create at least a block for the cytotoxin and tame it's sulphurous action of decay and render the

hit as merely haemotoxic and thence easier to handle pain-wise, facilitating a detox with **less** amounts of the iron antitoxin:..."*open bottle – drink a few sips – keep still!*"....*call a helicopter if you can*. The iron will convert to caffeine in the blood stream so that you remain **AWAKE!** We cannot guarantee that you will survive a cytotoxic attack without losing a limb, but at least you may have chances of staying alive. Always listen to local bushmen who may have useful tips about these things, - and don't play the worldly-wise dare-devil.

Long Storage Time:

Suited as a one-time long-term investment, snake antidote can also be stored for:

- **3 years without refrigeration in it's medical plastic bottle**, in a cool dry place away from sunlight and hot heat sources, secure it away from plastic-gnawing rodents or tropical termites and ants that are known to eat silicon plastic, cable coatings, plastic packaging and synthetic utensils that are exposed to salty coastline ozone.
- **50 years in refrigeration**, - but use at room temperature (do not deep freeze! Do not store in a cooler box!).
- **100 years if transferred into a glass bottle**, the antidote will remain potent, although glass bottles are heavy and very unsafe for luggage in the outback.

Precautions:

Always read the user instructions and package leaflet about amounts before using any biological detox and before possible exposure to toxic danger. **Wrong use other than detailed in the product's user-instructions does not legally constitute our liability.** The antitoxin is not recommended to replace a balanced diet, nor is it a lifestyle drug. For the treatment of minor complaints we recommend that users consult a trusted physician or doctor before considering any dosage plan in our detox application prior to purchase. Ascertain from your physician if a minor detox can be otherwise undertaken by changing diet habits, if needing to balance costs to upkeep a healthy diet. This antidote can be used by overweight diabetics under close medical supervision, though this detox is not recommended for slim or underweight diabetics. For further dosage details see the amount recommendations below.

Contraception vs. detox:

Modern everyday chemical birth contraception are often outlawed in many tropical nations for vital reasons, because they complicate all detox treatments against tropical illness in hot climates. The necessary high levels of formalin plasticity which chemical contraception employs, would usually facilitate the prevention of unplanned pregnancy in colder temperate Western or European climates, but can conversely result in life endangering thermic complications during any professional treatment of a tropical infection in southern or eastern tropical zones. Using chemical

means of contraception such as IUDs, spiral implants or contraceptive pills in hot tropical climates will complicate or even compromise the efficacy of all safe medical detox treatments in tropical areas. The bio-chemical reason for this, is that the high protein-plasticity barrier that makes contraception work, will prevent all life-saving detox medication from being effective, especially when high amounts of water intake during fever may further dilute and lower vital haemoglobin count, only to raise insubordinate white blood-cell levels. It is therefore strongly advised that women who contemplate travelling into tropical hot regions, should interrupt or replace such a chemical contraception plan with non-chemical means of contraception (i.e. *condoms, female condom or diaphragm*), until they return back home to the temperate zones where a high plasticity barrier has better chances of withstanding high body temperatures and exposure to sunshine. If cervical ripening during intercourse compromises condoms or diaphragm injury occurs, then temporary vas-occlusive contraceptive methods may be considered. Any treatment of Malaria or toxic illness can only be successful in the country of destination if chemistry is kept at normal level in the body. In such a case, snake antidote will prove a useful remedy for treating any toxicity arising from unsafe drinking water, suspicious meals or potentially toxic bites and street drugs.

Detox and Sexual Function vs. Dysfunction:

This antitoxin should not be used as a fertility measure if you or your partner is underweight or barren - and neither may this antitoxin prevent pregnancy at such a time. **Underweight males and females** may lose sexual interest if they use an overdose of the antitoxin when not necessarily treating envenomation or illness, reducing sexual libido and performance. However, in the case of a snake bite, it is usual that victims slowly recovering from the deadly malaise will experience a severe sexual urge during recovery, as poison toxins will convert to usable trace elements or cross-over cell-hormones in the body. The height and weight of a person's anatomy doesn't always correspond to an illness that they may have and reactions will vary from person to person. Underweight individuals do bounce back faster from toxic affliction after using less antidote, than normal-weight individuals per se, because their metabolic rate has less weight loss to manage before the treatment is over. In **regular-weight persons**, the antitoxin will raise sexual libido and performance in guarded doses, yet full amounts of antitoxin may be required to treat envenomation. Therefore, the use should be limited to a small amount if merely suffering iron deficiency from malnutrition (see amount instructions below). Compared to natural iron-rich diet, the boost of an iron-based antitoxin in level doses acts as a body-toning tonic that can raise sexual libido only in people with a normal diet or normal-to-heavy bodyweight.

Pregnancy:

Pregnant mothers should seek added necessary advice before using this or any non-toxic detox or follow the amounts states below. This antitoxin should not be used as a fertility measure if you or your partner is underweight or barren - and neither may this antitoxin prevent pregnancy at such a

time. Underweight along with depleted protein levels will most certainly reduce desired sexual libido and hinder both sperm count and egg cell production if this high-iron detox is used, hence forfeiting it's intended purpose. Chemical birth contraception may be compromised when using any detox, therefore prior consultation with a gynaecologist is highly advised...and better yet, your toxicologist.

Children:

Before using this antidote on a child, paediatric advice ought to be sought about the amounts of iron that a child may need according to it's weight and eating habits, if the antidote's instructions [here](#) are unclear to you. When not treating a severe toxic emergency in children, it is wise to keep the tone of tonics at small doses described in the user instructions, because children are usually very active at play and have higher metabolic rates compared to adults, because their little bodies cover more distance than grown adults do - especially while playing in warm climates. Too much antitoxin frequently administered to a child can stunt a child's growth if it's diet is wrong, while the correct dose and balanced meals can promote an even bone structure, stronger teeth and mental clarity. When there is no chronic illness, less tonic in the size of drips often does more good to a child's growth than liberal doses. Young people overcome sickness faster, largely due to their active nature and their propensity to consume needed sweets and fats during growth spasms. Skinny or less greedy children often tend to be more reclusive and sickly than their loud boisterous peers, though becoming great thinkers in the process. **So, as sad as it sounds, keep in mind that the smaller size of a child's body may immediately succumb to any fully virotoxic attack as compared to an adolescent or grown adult, - regardless of the amount of antidote administered afterwards.** Here is where prevention is better than cure. So, take this advice: It is reassuring to know that a child can survive a poisoning if a full dose of antitoxin is applied in the very instant that a fully toxic attack has become apparent. Saving a small child from a snake bite is very labour intensive and usually calls for hospitalization. Their small bodies cannot contain or pocket expulsive material well. All the more imperative it is, to also ply a child with steroid builders (i.e. calcium, magnesium or biotin) after a serious illness has been overcome. Moreover, a correct protein rich nutrition after any detox in a child will reward the effort. This is why it is wise to try to keep your children out of harm's way. One method of protecting children who frequently encounter poisonous wildlife is to keep an antitoxin in the medicine cabinet and out of their idle reach.

Recommended Dosages by Graph in Kilograms:

The divine rule of toxicity in bio-chemistry teaches: *"Too much of any good thing is always BAD for you!"*. Whether you gorge yourself on ice-cream or just on salads, you will learn that the exclusion of one thing above another is like racism in the chemistry of your body.

User guide below:

PATIENT'S ADVISORY**SNAKE ANTIDOTE****READ BEFORE USE!****WWW.SNAKE-ANTIDOTE.COM***Intl. Product Patent Nr. RT 20 603 031 5DE 200***500 ml / Type: SAFARI BIO-MEDICINE - SNAKE ANTIDOTE
100% Botanic / or + Capsiacin**

To be taken as follows if not otherwise directed by our designated health physicians:

QUICK EMERGENCY USE AT A GLANCE

Read carefully and pay attention to age, application, and **quantity** amounts below. **Drink directly from the bottle after snake bite!** Do not take any other medications with any snake antidote. Do not dilute! Avoid alcohol, meat-diet, fish, cheese or poultry with this antitoxin!

The speed application after a snake bite is intended to treat toxic **Hyper-Insulin-Volumes (HIV-complex)** within 24 hours! Snakes, poisonous reptiles and insects vary! It is difficult to know in advance how much poison a snake has injected with its sharp fangs. Likewise it is equally difficult for any doctor to know all the risky things a patient may have done before becoming ill. Therefore, we start a treatment with Snake Antidote as a classic purging process to diminish toxic viro-proteins.

SIDE EFFECTS

This antidote will cause violent necessary toilet expulsion of poisons, dripping nose, and temporary high blood pressure, occasional dizzy spells, and odd nocturnal dreams. Before taking it, you may prime stomach with half glass of salt water. For preliminary questions write to the email address in our company address below at the foot of this instruction.

SNAKE BITE: QUANTITY / DOSESMature snake bite = **2-4 bottles** with speed - all in.Juvenile snake bite = **1-2 bottles with speed** - all in.

The rule is to administer full units until the patient feels fine and is **consciously** out of clinical danger. **For all other less life threatening infections Children and infants may take guarded strongly reduced amounts (see graph below!).** Infants under 2 years are difficult to save after a full snake bite, but may be easily cured of other Hyper Insulin Volumes (HIV-complex) such as Hepatitis and Leukaemia that can come from Malaria affliction.

All Other Infections			
AGE	AMOUNT	DURATION	MODE
Infant till 11 months	2 x 10 ml	1 x week / max. 2 weeks	1 nursing-bottle with sugar and milk
Baby 1 year old	2 x 25 ml	1 x week / max. 2 weeks	1 nursing-bottle with sugar and milk
Baby 2 year old	2 x 30 ml	1 x week / max. 2 weeks	1 nursing-bottle with sugar and milk
Toddler 3 year old	2 x 50 ml	1 x week / max. 2 weeks	1 nursing-bottle with sugar and milk
Child 4 - 6 years old	2 x 50 ml	2 x week / max. 2 weeks	unadulterated
Child 7 - 10 years old	2 x 60 ml	2 x week / max. 2 weeks	unadulterated
Pre-Teen 11 - 15 y-old	2 x 60 ml	2 x week / max. 2 weeks	unadulterated
Pubescent 16 – 18 y	2 x 100 ml	1 x week / max. 2 weeks	unadulterated
Adolescent 20 – 27 y	2 x 100 ml	1 x week / max. 2 weeks	unadulterated
Adult 28 and upwards	1 half-bottle 120 ml	2 x week / max. 2 weeks	unadulterated

RESPONSE: Body responds within 2 – 5 minutes. The patient will feel extreme revival.

Discontinue use if treatment induces an imbalanced protein deficiency or underweight.

PREGNANT MOTHERS			
STAGE	AMOUNT	DURATION	MODE
Healthy pregnant mothers under 3 months	3 drops a day	2 weeks	with milk + normal diet incl. meat, poultry, fish.
HIV diagnosed pregnant mothers under 3 months	1 x 25 ml a week	2 weeks	with milk + normal diet incl. meat, poultry, fish.
RESPONSE: Body responds within 2 – 5 minutes. The patient will feel extreme revival.			

Discontinue use if treatment induces an imbalanced protein deficiency or underweight.

SAME TREATMENT PLAN FOR:

MALARIA STRAINS & WEST NILE VIRUS
 OVERWEIGHT DIABETES (**not for underweight Diabetics!**)
 HEPATITIS
 LEUKAEMIA
 SICKLE-CELL ANAEMIA & ANAEMIA (iron deficiency)
 1st STAGE CANCER (Metastasis)
 PALLIATIVE DRUG ADDICTION
 OBESITY
 TOXIC CARCINOGENIC POISONING
 SEX-LUBRICANT OVERDOSE (Vaginal / Anal)

Discontinue use if treatment induces an imbalanced protein deficiency or underweight.

This antidote may be taken while smoking an emergency cigarette or cubana for neural relaxation, but can aid in curbing the smoking habit. The antidote works most effectively on smokers or people who eat salt, but slower on non-smokers and people who abstain from salt. Do not use nicotine patches or nicotine sweets during a detox.

OBESITY & OVERWEIGHT DIABETES TREATMENT
 (Not for slim Diabetics)

OVERWEIGHT DIABETICS			
WEIGHT LOSS	AMOUNT	DURATION	MODE
3 kg	1 antidote unit 50ml 5 x a week	2 weeks or extendable over months per kg	Rich vegan high carbohydrate diet with very sweet fruit beverages and 1 magnesium pill at 600mg a day
CALCULATION: According to desired kg loss until average weight x height			

Discontinue use if treatment induces an imbalanced protein deficiency or underweight.

WARNING: ADVICE FOR OVERWEIGHT DIABETICS:

Overweight Diabetics don't know that most of their weight-gain is 50% oedema and 50% clogged fats. However, diabetics do know that their CE levels (meaning: "carbohydrate exchange") are necessary to evaluate how much carbohydrates such as in bread or barley will convert to **insulin** in the pancreas, but are afraid of getting **fat** (*insulin*) from the same strong insulin source (bread) which actually keeps an acidic lack on insulin at bay. **So, the idea of using an iron-based antitoxin to transform overt toxic stagnant fats and cholesterol into lots of needed fluid insulin, means that in the instant that a diabetic uses a dark antitoxin to liquefy stagnant fats, they aren't diabetics any more. This phenomenon happens within 3 days of using snake antitoxin.**

This favourable condition continues for the entire duration of the **safe** weight-loss plan of **using 50ml antitoxin 5 x a week for 2 weeks on a highly carbohydrate rich vegan diet with sweet fruit beverages and 1 magnesium pill at 600mg a day as a diuretic**. The plan can extend beyond two weeks for all the months that an obese person needs to lose weight. The iron-based antitoxin allows diabetics who previously were unable to eat white bread, to digest white wheat without any liver complaints.

Diabetes may return however, right after an optimal slimline body is rapidly achieved if they didn't grasp the role that correct amounts of carbohydrates play in dietary insulin supply. **Using another antidote program will not work on fresh slim diabetics**. Formerly fat diabetics are capable of learning, that during liquid fat mobilisation, all the needed insulin was trapped for years as stagnant fat deposits around the belly all along because of a false understanding that FAT is **insulin** that needs even distribution away from just the belly, if sweet beverages transport it.

Therefore, high amounts of CE-carbohydrate-exchange levels and normal fruit beverages cause optimal insulin intake and level weight management, when newly trim diabetics return to eating meats and other similar proteins with sweet fruit beverages after the weight-loss plan is over. Why? Meat and other animal proteins and even steroid protein precursors such as a magnesium, biotin and calcium will convert to added insulin with the use of sweet beverages with every oily meal containing salt.

Diabetics must LEARN during weight loss how vital the use of SUGAR is for the heart-rate while using a bitter detox of fats, or else they will suffer cardiac arrest from oedema and liquid fats filling the pericardium, precisely because a lack of SUGAR causes **hypoglycaemia**. Most diabetics aren't prepared to accept this toxicological fact about low glucose levels, reasoning that their bodies are different than everybody else's. Though SUGAR promotes excellent skin elasticity during rapid weight loss needed to prevent undulating flabby skin bags and hair loss after losing the bulge, many diabetics are mortally afraid of sugar.

Also, Diabetics who are on hydrochloric pills (Metformin or similar) while also taking insulin injections should note that hydrochloric pills reduce the efficacy of insulin injections and also bring the vital haemoglobin stats down, causing bouts of **diabetic anaemia**. They occasionally grab a glass of orange juice put next to the bed at night, while many healthy and slim individuals drink an entire bottle after eating a fatty steak! Diabetics don't know that drinking fruit beverages while eating protein creates insulin and weight loss. These were people who never understood the sexy relationship that sugar has with beef.

Because many diabetics become fearfully phobic to all sugar and fruits they begin to become overweight by drinking mainly bland water which again flushes away healthy iron levels that level sugar counts provide, which leads to more weight gain and water retention. When WATER further dilutes the iron count, it creates the namesake Greek description *DIABETES* (meaning: **blanched red**) associated with **low blood sugar (the iron precursor)**.

In addition, the use of cigarettes, black coffee, dark greens or any dark meal without sugar can trigger a false high sugar reading even though the insulin levels are stable – but these false sugar levels will subside usually some **30 minutes after digesting such a meal**. - Most diabetics don't wait to see the sugar levels dip naturally and hasten to inject more insulin which jacks up the white blood cell differential, then grab a bottle of water when the throat begins to itch. Oedema rises! **Likewise, when using this iron-based detox to shed unwanted weight gain, bitter iron can trick your glucometer into telling you to inject insulin prematurely, which in turn causes you to suddenly develop jaundice, which in effect is an insulin overdose! Even though an insulin overdose does not apply to diabetics and doesn't lead to ketoacidosis, but can lead to heart failure simply because the calming sanguinity of blood-sugar is missing. Although this complication in treating iron deficiency in diabetics calls for that famous life-saving drink of orange juice standing next to the bed, - many diabetics refuse to even touch a drink of fruit juice**. Stubbornly, they go ahead to drink lots of great-tasting commercial structured bleach-water or even down a salty meal, refuting the weight-loss plan they begged for, only to enter a state of **leukaemia**. The fact that dark bitter iron tonics render salty body fats “*yellow*” when SUGAR is missing, means that a diabetic can go from being insulin deficient to having a false HIV-complex, which can cause a **false sexual stigma**.

Diabetics learn that in the instant that they become leukaemic from shedding liquefied unwanted FAT, they will have to eat sweets like any victim of Malaria to expel those fats without losing HAIR and TEETH at the same time. Therefore, we distance ourselves from any wrong diabetic behaviour, if they don't respect the laws of toxicology that Diabetologists convey when teaching their patients about CE-

carbohydrate exchange levels necessary to evaluate how much dark bread or other carbohydrates convert to insulin, for a glass of orange juice to save them from hypoglycaemia!

If contemplating using this detox to convert overweight fat into healthy insulin levels, we welcome that diabetics who wish to lose weight by converting their excessive body fat and oedema to insulin, consult with their diabetologists who may accompany their quest for a better understanding of seroconversion.

Understanding that a carbohydrate rich vegan diet increases healthy weight loss while keeping correct insulin levels with the use of sugar, can improve their quality of life.

LIVESTOCK

PETS & FARM-ANIMAL SUPPLEMENTATION / By Minor Average Values					
SPECIMEN	Kg	Lbs	AMOUNT	DURATION	MODE
Poultry / Avian	1.0 – 3 kg	2.2 lb	2 x 10 ml	1 x week / max. 2 weeks	feed kaolin+sugar
Mouse	0.02 kg	0.04 lb	2 x 4 ml	1 x week / max. 2 weeks	feed-gun / feed
Hamster	0.12 kg	0.26 lb	2 x 4 ml	1 x week / max. 2 weeks	feed-gun / feed
Guinea pig	1.04 kg	3.0 lb	2 x 4 ml	1 x week / max. 2 weeks	feed-gun / feed
Rabbit	2.5 kg	5.5 lb	2 x 10 ml	1 x week / max. 2 weeks	feed-gun / feed
Cat	3.3 kg	7.2lb	2 x 10 ml	1 x week / max. 2 weeks	feed-gun / feed
Dog	14 – 25 kg	30 – 55 lb	2 x 10 ml	1 x week / max. 2 weeks	feed-gun / feed
Lamb / Goat	18.1 – 63.5 kg	40 – 140 lb	2 x 10 ml	1 x week / max. 2 weeks	feed-gun / feed
Pig	5.5 – 12.5 kg	12 - 34 lb	2 x 10 ml	1 x week / max. 2 weeks	feed-gun / feed
Calf	25 – 45 kg	55 - 99 lb	2 x 50 ml	1 x week / max. 2 weeks	feed-gun / feed
Cattle / Horses	259 – 800 kg	551–1800 lb	2 x 100 ml	1 x week / max. 2 weeks	feed-gun / feed
For other species gage according to weight relation listed.					
RESPONSE: Body responds within 2 – 5 minutes. The patient will feel extreme revival.					

Discontinue use if treatment induces an imbalanced protein deficiency or underweight.

ADDED USE ON LIVESTOCK

The snake antitoxin is not just used to treat poisonous snake or scorpion bites, but also can be used against the following diseases to bring down fevers and malaise. It is introduced as a purger to a sick animal's mouth via a pet feeding gun or mouth-syringe or simply mixed into the feed with a little bit of sugar, which will do the animal a world of good:

Used for:

Anaplasmosis
 FMD (Foot & Mouth Disease)
 Bovine Respiratory Disease (BRD)
 Blackleg
 Johne's Disease
 Bovine Tuberculosis (TB)

Not for:

Bovine Viral Diarrhea (BVD) / use kaolin-pectin preparations or electrolyte solutions
 Mastitis / use kaolin-pectin preparations or electrolyte solutions

CATS & DOGS

Other than against meat poisoning, this antitoxin is given only to dogs and cats in the winter time, when the natural young grass they seek for bile acids are hard to find to clean their spleen.